

Atomic Energy Central School No. 4, Rawatbhata

Class XII (Physics, Chemistry, Mathematics/Biology)

Multiple Choice Questions Examination - July (2019-20)

Name of student: _____ Class: _____ Roll No. _____

General Instructions: 1. Darken the appropriate circle in the OMR answer sheet.


2. Each question carries 1 mark. There is no negative marking.

Physics

1. A half ring of radius R has a charge of λ per unit length. The field at the center is


- a) $2 \frac{k\lambda}{R}$ b) $\frac{k\lambda}{R}$
c) zero d) $\frac{n\lambda}{R}$

2. A uniformly charged thin spherical shell of radius R carries uniform surface charge density of σ per unit area. It is made of two hemispherical shells, held together by pressing them with force F (See figure). F is proportional to




- a) $\frac{\sigma^2 R^2}{\epsilon_0}$ b) $\frac{\sigma^2 R}{\epsilon_0}$
c) $\frac{1}{\epsilon_0} \frac{\sigma^2}{R^3}$ d) $\frac{1}{\epsilon_0} \frac{\sigma^2}{R}$

3. Consider a system of three charges $\frac{q}{3}$, $\frac{q}{3}$ and $-\frac{2q}{3}$ placed at points A, B and C, respectively, as shown in the figure. Take O to be the centre of the circle of radius R and angle $CAB = 60^\circ$



- a) The electric field at point O is $\frac{q}{4\pi\epsilon_0 R^2}$ directed along the negative x-axis b) The magnitude of the force between the charges at C and B is $\frac{q^2}{54\pi\epsilon_0 R^2}$
- c) The potential energy of the system is zero d) The potential at point O is $\frac{q}{12\pi\epsilon_0 R}$



a) $\frac{q}{2\pi\epsilon_0 L}$

b) None of these

c) $\frac{q}{4\pi\epsilon_0 L}$

d) $\frac{q}{3\pi\epsilon_0 L}$

11. Consider a uniform electric field $E = 3 \times 10^3 \text{ N/C}$. 1

- a. What is the flux of this field through a square of 10 cm on a side whose plane is parallel to the yz plane?
- b. What is the flux through the same square if the normal to its plane makes a 60° angle with the x-axis?

a) $30 \text{ Nm}^2/\text{C}$, $15 \text{ Nm}^2/\text{C}$

b) $20 \text{ Nm}^2/\text{C}$, $15 \text{ Nm}^2/\text{C}$

c) $40 \text{ Nm}^2/\text{C}$, $15 \text{ Nm}^2/\text{C}$

d) $40 \text{ Nm}^2/\text{C}$, $25 \text{ Nm}^2/\text{C}$

12. Careful measurement of the electric field at the surface of a black box indicates that the net outward flux through the surface of the box is $8.0 \times 10^3 \text{ Nm}^2/\text{C}$. 1

1. What is the net charge inside the box?
2. If the net outward flux through the surface of the box were zero, could you conclude that there were no charges inside the box?

a) $0.04 \mu\text{C}$, Yes

b) $0.06 \mu\text{C}$, Yes

c) $0.05 \mu\text{C}$, No

d) $0.07 \mu\text{C}$, No

13. A force of repulsion between two point charges is F, when these are at a distance 0.1 m apart. Now the point charges are replaced by conducting spheres of radii 5 cm each having the same charge as that of the respective point charges. The distance between their centres is again kept 0.1 m, then the force of repulsion will: 1

a) remain F

b) decrease

c) increase

d) become $\frac{10F}{9}$

14. A semi-circular arc of radius 'a' is charged uniformly and the charge per unit lengths is λ . The electric field at the centre is: 1

a) $\frac{\lambda}{2\pi\epsilon_0 a^2}$

b) $\frac{\lambda}{4\pi\epsilon_0 a}$

c) $\frac{\lambda}{2\pi\epsilon_0 a}$

d) $\frac{\lambda^2}{2\pi\epsilon_0 a}$

15. A metal plate of thickness half the separation between the capacitor plates of capacitance C, is inserted between the plates. The new capacitance is 1

a) $\frac{c}{2}$

b) 0.0

c) C

d) 2C

16. To make a condenser of $16\mu\text{F}$, 1000 volts, how many condensers are needed which have written on them " $8\mu\text{F}$, 250 volts"?

a) 8.0

b) 32.0

c) 40.0

d) 2.0

17. Two identical capacitors, have the same capacitance C. One of them is charged to potential V_1 and the other to V_2 . The negative ends of the capacitors are connected together. When the positive ends are also connected, the decrease in energy of the combined system is

a) $\frac{1}{4}C(V_1 - V_2)^2$

b) $\frac{1}{4}C(V_1^2 + V_2^2)$

c) $\frac{1}{4}C(V_1^2 - V_2^2)$

d) $\frac{1}{4}C(V_1 + V_2)^2$

18. A $2\mu\text{F}$ capacitor C_1 is charged to a voltage 100 V and a $4\mu\text{F}$ capacitor C_2 is charged to a voltage 50 V. The capacitors are then connected in parallel. What is the loss of energy due to parallel connection?


a) $1.7 \times 10^{-2} \text{J}$

b) $0.17 \times 10^{-2} \text{J}$

c) 1.7 J

d) $1.7 \times 10^{-4} \text{J}$

19. Three capacitors, each of capacitance $C = 3 \text{ mF}$, are connected as shown in the figure. The equivalent capacitance between points P and S is



a) $3 \mu\text{F}$

b) $9 \mu\text{F}$

c) $1 \mu\text{F}$

d) $6 \mu\text{F}$

20. A variable capacitor and an electroscope are connected in parallel to a battery. The reading of the electroscope would be decreased by


a) Decreasing the battery potential

b) Increasing the area of overlapping of the plates

c) Decreasing the distance between the plates

d) Placing a dielectric between the plates


21. A parallel plate air filled capacitor shown in the Fig. (a) has a capacitance of $2 \mu\text{F}$. When it is half filled with a dielectric of dielectric constant $k = 3$ as shown in Fig. (b), its capacitance becomes



- a) $\frac{1}{3} \mu\text{F}$ b) $9 \mu\text{F}$
 c) $1 \mu\text{F}$ d) $3 \mu\text{F}$

22. Two capacitors A and B are connected in series with a battery as shown in figure. When the switch S is closed and the two capacitors get charged fully then

1



- a) the potential difference across the plates of A is 4 V and across the plates B is 6 V b) the ratio of charges on A and B is 3 : 2
 c) the ratio of electrical energies stored in A and B is 2 : 3 d) the potential difference across the plates of A is 6 V and across the plates of B is 4 V


23. A parallel plate capacitor of value $1.77 \mu\text{F}$ is to be designed using a dielectric material (dielectric constant 200, breakdown strength of $3 \times 10^{-6} \text{Vm}^{-1}$). In order to make such a capacitor, which can withstand a potential difference of 20 V across the plates, the separation d between the plates and the area A of the plates should be

1

- a) $d = 10^{-5}\text{m}$, $A = 10^{-2}\text{m}^2$ b) $d = 10^{-4}\text{m}$, $A = 10^{-4}\text{m}^2$
 c) $d = 10^{-4}\text{m}$, $A = 10^{-5}\text{m}^2$ d) $d = 10^{-6} \text{m}$ and $A = 10^{-4}\text{m}^2$

24. A parallel plate air filled capacitor shown in Fig. (a) has a capacitance of $2 \mu\text{F}$. When it is half filled with a dielectric of dielectric constant $k= 3$ as shown in Fig. (b), its capacitance becomes

1



a) $0.5 \mu\text{F}$

b) $3 \mu\text{F}$

c) $4 \mu\text{F}$

d) $1.5 \mu\text{F}$

25. A parallel plate capacitor of plate area A has a charge Q . The force on each plate of the capacitor is **1**

a) $\frac{2q^2}{\epsilon_0 A}$

b) zero

c) $\frac{q^2}{\epsilon_0 A}$

d) $\frac{q^2}{2\epsilon_0 A}$

26. If the electric current in a lamp decreases by 5%, then the power output decreases by: **1**

a) 20%

b) 25%

c) 10%

d) 5%

27. A potentiometer has a uniform wire of length 10m and resistance 5 ohms. The potentiometer is connected to an external battery of emf of 10V and negligible internal resistance and a resistance of 995 ohms in series. The potential gradient along the wire is: **1**

a) 1 mV/cm

b) 5 mV/cm

c) 1 mV/m

d) 5 mV/m

28. Power dissipated in a resistance R through which current I is flowing is **1**

a) $I^2 R$

b) $I^2 R^2$

c) IR

d) IR^2

29. According to Ohm's law **1**

a) The electric current I flowing through a substance is proportional to the voltage V across its ends

b) The electric current I flowing through a substance is proportional to the square of voltage V across its ends

c) The electric current I flowing through a substance is inversely proportional to the voltage V across its ends

d) The electric current I flowing through a substance is independent of the voltage V across its ends

30. An electric kettle taking 3 A to 200 V brings one litre of water from 20°C to the boiling point in 10 minute. Its efficiency is: **1**

a) 93.0%

b) 33.3%

c) 66.6%

d) 87.7%

31. The resistance of a metallic conductor increases due to **1**

a) Change in dimensions of the conductor

b) Change in carrier density

c) Increase in the number of collisions between the carriers

d) Increase in the rate of collisions between the carriers and vibrating atoms of the conductor

32. Which can be the units of Resistivity?

1

a) $meter \times \frac{Ampere}{Volt}$

b) $Volt \times \frac{Ampere}{meter}$

c) $\frac{Volt \ meter}{Ampere}$

d) $Volt \times Ampere$

33. The wire of the potentiometer has resistance 4 ohms and length 1 m. It is connected to a cell of e.m.f. 2 volts and internal resistance 1 ohm. The current flowing in the potentiometer is:

1

a) 0.4 A

b) 0.1 A

c) 0.8 A

d) 0.2 A

34. Current density of a conductor is

1

a) Is always zero

b) the net charge flowing through the area

c) the net current flowing through the area normally per unit time

d) the net charge flowing through the area per unit time

35. Direction of the conventional current

1

a) is the direction in which negative charges move

b) is the direction in which positive charges move

c) is the direction in which no charges move

d) to the direction in which positive charges move

36. Orders of magnitude of random electron motion speed to drift speed are like

1

a) $10^2 \text{ m/s}, 10^2 \text{ m/s}$

b) $10^3 \text{ m/s}, 10^{-1} \text{ m/s}$

c) $10^4 \text{ m/s}, 10^{-2} \text{ m/s}$

d) $10^6 \text{ m/s}, 10^{-4} \text{ m/s}$

37. An ammeter together with an unknown resistance in series is connected

1

across two identical batteries each of emf 1.5 V. When the batteries are connected in series, the galvanometer records a current of 1A and when the batteries are in parallel, the current is 0.6A. What is the internal resistance of each battery?

a) $\frac{1}{5} \Omega$

b) $\frac{1}{3} \Omega$

c) $\frac{1}{4} \Omega$

d) $\frac{1}{2} \Omega$


38. According to Kirchhoff's Loop Rule

1

- | | |
|---|---|
| a) The absolute sum of changes in potential around any closed loop must be zero. | b) The algebraic sum of changes in potential around any closed loop must be zero. |
| c) The algebraic sum of changes in potential around any closed loop must be positive. | d) The algebraic sum of changes in potential around any closed loop must be negative. |

39. In the circuit shown below, the cell is ideal, with emf = 2 V. The resistance of the coil of the galvanometer G is 1Ω

1



- | | |
|--|--|
| a) 0.2 A current flows in G. | b) Potential difference across C_2 is 1.2 V. |
| c) Potential difference across C_1 is 1 V. | d) No current flows in G. |

40. If the number of turns, area and current through a coil is given by n , A and I respectively, then its magnetic moment will be:

1

- | | |
|------------------|------------|
| a) nI/\sqrt{A} | b) nIA |
| c) n^2IA | d) nIA^2 |

Chemistry

41. The half life periods of a reaction at initial concentration 0.1 mol/L and 0.5 mol/L are 200 s and 40 s respectively. The order of the reaction is

1

- | | |
|------|------------------|
| a) 2 | b) $\frac{1}{2}$ |
| c) 0 | d) 1 |

42. For an endothermic reaction where ΔH represents the enthalpy of the reaction in kJ/mol. The minimum value for the energy of activation will be

1

- | | |
|-------------------------|-------------------------|
| a) Equal to ΔH | b) Zero |
| c) More than ΔH | d) Less than ΔH |

43. If a reaction proceeds with a uniform rate throughout, the reaction is

1

- | | |
|----------------|-----------------|
| a) Third order | b) Second order |
| c) First order | d) Zero order |


44. Rate of reaction does not remain constant throughout because 1
- a) Density of reactants keep on changing b) Concentration of reactants keep on changing
- c) Volume of reactants keep on changing d) Temperature of reactants keep on changing
45. The rate law for the reaction is given by rate = $k[RCl]$. The rate for this reaction 1
- a) is unaffected by change in temperature b) is halved by doubling the concentration of NaOH
- c) is doubled by doubling the concentration of NaOH d) is halved by half by reducing the concentration of RCl
46. Thermal decomposition of a compound is of first order. If 50% of a sample of a compound is decomposed in 120 min, the time taken for 99.9% completion is 1
- a) 1000 min b) 399 min
- c) 1200 min d) 400 min
47. The slope in the $\log k$ vs $\frac{1}{T}$ curve is 5.42×10^3 . The value of the activation energy is approximately
- a) 104 J/mol b) 208 J/mol
- c) 104 kJ/mol d) 104 J/mol
48. The reaction $A + 2B \rightarrow C + D$ obeys the rate equation Rate = $k[A]^x[B]^y$ what would be the order of this reaction? 1
- a) x b) x + y
- c) x - y d) Cannot be predicted with the equation
49. Which among the following statement is not true for rate constant of a reaction? 1
- a) Unit of rate constant depend upon the order of reaction b) Rate constant depend upon the concentration of the reactants
- c) Rate constant has a definite value at a particular temperature d) Rate constant changes with temperature
50. The reaction $2NO + Br_2 \rightarrow 2NOBr$ follows the mechanism given below 1
- $NO + Br_2 \rightleftharpoons NOBr_2$ (fast)
- $NOBr_2 + NO \rightarrow 2NOBr$ (slow)
- If the concentration of both NO and Br_2 are increased two times, the rate of reaction would become
- a) 2 times b) 8 times
- c) 4 times d) 6 times

51. The units for the rate constant for the second order reaction (concentration : mol litre⁻¹ time: s) are: 1
- a) s⁻¹ b) mol litre⁻¹ s⁻¹
 c) mol litre⁻² s⁻¹ d) mol⁻¹litre s⁻¹
52. Reaction which takes place in one step is known as 1
- a) Elementary reaction b) Unimolecular reaction
 c) Reaction rate d) Bimolecular reaction
53. For a chemical reaction 2X + → Z, the rate of appearance of Z is 0.05 mol L⁻¹min⁻¹ 1
1. The rate of disappearance of X will be
- a) 0.05mol L⁻¹min⁻¹ b) 0.1 molL⁻¹min⁻¹
 c) 0.25mol L⁻¹min⁻¹ d) 0.05mol L⁻¹hour⁻¹
54. Which of the following rate laws is third order overall? 1
- a) rate = K[A]⁵[B]² b) rate = K [A] [B]²
 c) rate = K[A]³[B]³ d) rate = K[A]³[B]¹
55. Which catalyst is used in Haber's process? 1
- a) Molybdenum b) Iron
 c) Platinum d) Vanadium
56. Which of the following reaction gives a colloidal sol? 1
- a) Cu + CuCl₂ → Cu₂Cl₂ b)
2HNO₃ + 3H₂S → 3S + 4H₂O + 2NC
 c) d) MgCO₃ → MgO + CO₂
 2Na + 2H₂O → 2NaOH + H₂
57. An example of autocatalysis is 1
- a) Decomposition of KClO₃ to KCl and O₂ b) Oxidation of NO to NO₂
 c) Oxidation of SO₂ to SO₃ d) Oxidation of oxalic acid by acidified KMnO₄
58. Which adsorption takes place at low temperature? 1
- a) Chemical b) Can not say
 c) Physical d) Both Physical and Chemical
59. The path of light becomes visible when it is passed through As S sol in water. 1
- (give reason)
- a) Due to Brownian movement b) Due to micelle formation
 c) Due to colour formation d) Due to Tyndall effect

60. Which is correct in case of Van der waal adsorption? **1**
- a) High temperature, high pressure b) Low temperature, high pressure
- c) Low temperature, low pressure d) High temperature, low pressure
61. The adsorbent used to adsorb the dye particles in the dyeing industry is **1**
- a) Activated charcoal b) Silica gel
- c) Alum d) Alumina gel
62. Which type of a property is the Brownian movement of colloidal solution? **1**
- a) Electrochemical b) Optical
- c) Mechanical d) Electrical
63. Micelles are: **1**
- a) Ideal solution b) Associated colloids
- c) Adsorbed solution d) Emulsion cum gel
64. Fog is a colloidal solution of **1**
- a) Liquid in gas b) Gas in liquid
- c) Solid in gas d) Gas in gas
65. Which of the following is not exhibited by solutions? **1**
- a) Absorption b) Flocculation
- c) Paramagnetism d) Tyndall effect
66. Which catalyst is used in contact process? **1**
- a) Molybdenum b) Vanadium pentoxide
- c) Platinum d) Iron
67. Which of the following processes does not involve a catalyst? **1**
- a) Thermite process b) Haber process
- c) Oswald process d) Contact process
68. In blast furnace, the highest temperature is in **1**
- a) Reduction zone b) Slag zone
- c) Fusion zone d) Combustion zone
69. Which among the following is a chemical process? **1**
- a) Magnetic separation b) Froth floatation
- c) Gravity separation d) Leaching
70. The cyanide process is used for obtaining **1**
- a) Ag b) Cu
- c) Zn d) Na

Biology


81. Given below is a highly simplified representation of human sex chromosomes 1 from a karyotype. The genes a and b could be of:



- a) Colour blindness and body height
b) Attached ear lobe and Rh blood group
c) Phenylketonuria and haemophilia.
d) Haemophilia and red green colour blindness
82. Which of the following is not a Mendelian disorder? **1**

- a) Hemophilia
b) Down's syndrome
c) Thalassemia
d) Colour blindness

83. Represented below is the inheritance pattern of a certain type of trait in humans. Which one of the following conditions could be an example of this pattern? **1**



- a) Thalassemia
b) Haemophilia
c) Sickle Cell anemia
d) Phenyl ketonuria
84. In somatic cells of human beings chromosomes exists as **1**
- a) Haploid
b) Heterologous pairs
c) Single circular chromosome
d) Homologous pairs

85. When two genes are situated very close to one another on a chromosome **1**

a) Hardly any cross-overs are produced
b) Only double cross-over can occur between them
c) No crossing over can take place
d) The percentage of crossing over between them is very high

86. Statement I: Physical association between two genes located on same chromosome is called linkage. **1**
Statement II: The generation of non-parental gene combination is called recombination.

Statement III: Recombination is essential for generation of new traits in a species.

- a) All statements are incorrect b) All statements are correct
 c) Only statement I and II is correct d) Only statement II and III are correct

87. A chromosome with sub-terminal centromere near the middle is called **1**

- a) Acentric b) Telocentric
 c) Acrocentric d) Metacentric

88. When two genes present on different loci produce the same effect when present alone but interact to form a new trait when present together are called **1**

- a) Polymeric genes b) Duplicate genes
 c) Complementary genes d) Supplementary genes

89. Failure of cytokinesis after telophase stage of cell division results in the **1**

- a) An increase in whole set of chromosome in organism b) Decrease in the whole set of chromosome in organism
 c) No change in the chromosome number d) Increase or decrease depends upon type of cell division

90. When chromosome sets are present in multiple of 'n' the condition is called **1**

- a) Haploidy b) Euploidy
 c) Aneuploidy d) Diploidy

91. Pedigree analysis is study of particular traits in **1**

- a) Two species b) Two families
 c) A several generation d) Two generation only

92. Whose karyotype is represented below? **1**



- a) Intersex Fly b) Male Drosophila
 c) Super female fly d) Female Drosophila

93. Henking X-body present in 50% of sperms are now known as **1**

- a) Lamp brush chromosomes b) Autosomes
 c) Y-chromosome d) X-chromosome


94. Map distance of genes on chromosome is calculated by 1

- a) Non-cross over percentage b) Recombination frequency of each gene locus
c) Cross over percentage d) Number of mutant genes

95. Which one of the following conditions in human is correctly matched with its chromosomal abnormality / linkage? 1

- a) Erythroblastosis foetalis - X – linked b) Down's syndrome - 44 autosomes +XXY
c) Colour blindness - Y – linked d) Klinefelter's syndrome - 44 autosomes +XXY

96. What does the chart give below represent? 1



- a) XX - XO type of sex determination b) XX - XY type of sex determination
c) XO - XX type of sex determination d) xy - xx type of sex determination

97. A phenomenon which works opposite to the linkage is 1

- a) Independent assortment b) Segregation
c) Mutation d) Crossing over

98. Hemophilia is more commonly seen in human males than in human females because 1

- a) This disease is due to a Y-linked recessive mutation b) A greater proportion of girls die in infancy
c) This disease is due to a X-linked recessive mutation d) This disease is due to a Y-linked dominant mutation

99. If a colour blind woman marries a normal vision man, their sons will be 1

- a) 3/4th colour blind and 1/4th normal b) 1/2 colour blind and 1/2 normal
c) All colour blind d) All normal visioned

100. In our society women are blamed for producing female children as the growth and development of child occurs inside the mother womb. This statement is 1

- a) sometimes correct b) Correct
c) incorrect d) Always correct

101. Statement I: DNA finger printing is highly reliable method of identification of individual involved in crimes. **1**

Statement II: DNA a fingerprinting is a sure method in solving paternity and maternity disputes.

Statement III: DNA fingerprinting can be used to cure HIV infection.

- a) All statements are correct b) Statement I and II is correct
c) Statement I and III is correct d) Statement II and III is correct

102. Autoradiogram of VNTR probe gives many band of different size. It differ from individual to individual except **1**

- a) Heterozygotic twins b) Real brothers
c) Monozygotic twins or identical twins d) Real sisters


103. In biochemical genetics the term gene is being replaced by **1**

- a) Anticodon b) Genome
c) Template d) Cistron

104. Typically DNA content of about 100000 cells or 1 microgram is required for fingerprinting. If the sample obtained is less it is increased by **1**

- a) Transcription of DNA in cells b) Elimination of DNA in cells
c) Translation of DNA in cells
d) Polymerase chain reaction (PCR) by amplification process

105. H-bonds between Cytosine and Guanine are **1**




- a) 1 b) 2
c) 3 d) 4

106. Transcription of DNA into mRNA occurs in the nucleus of the cell but translation occurs in **1**

- a) Cytoplasm b) Mitochondria
c) Golgi apparatus d) Nucleus

107. What does X represent in the following diagram:

1



- a) Released tertiary protein
- b) Released polypeptide chain
- c) Released secondary protein
- d) Released 3D protein molecule

108. Which one of the following techniques made it possible to genetically engineer living organisms?

1

- a) X-ray diffraction
- b) Hybridization
- c) Heavier isotope labeling
- d) Recombinant DNA techniques

109. Single nucleotide polymorphism (SNPs) revolutionize the process of finding chromosomal locations for

1

- a) Treatment of sex-linked genes
- b) Hybridization
- c) Disease-associated sequences and tracing human history
- d) Fingerprinting

110. Removal of introns and joining of exons in a defined order during transcription is called

1

- a) Splicing
- b) Inducing
- c) Looping
- d) Slicing

111. What is the purpose of the Human Genome Project?

1

- a) To archive everyone's DNA fingerprint
- b) To develop an "ideal" genetic code
- c) To mass produce important sequences of DNA
- d) To identify the sequence of human DNA

112. A gene of operon which synthesizes a repressor protein is

1

- a) Structural gene
- b) Regulator gene
- c) Operator gene
- d) Promoter gene

113. According to the lac-operon concept, which functional unit of the bacterial gene material is responsible for suppressing the activity of the operator gene in the absence of lactose?

1

- a) Repressor protein
- b) Structural gene
- c) Promoter gene
- d) Regulator gene

114. What does "lac" refer to in what we call the lac operon? **1**
 a) Lactase
 b) Lactose
 c) Lac insect
 d) The number 1,00,000
115. DNA probes used in finger printing are **1**
 a) Highly sensitive electron microscope
 b) UV beams
 c) X-ray scanners
 d) DNA segments having radioactive isotopes
116. Human genome project can leads to revolutionary new ways to **1**
 a) Study the mechanism of disease development
 b) Diagnose, treat and prevents the thousands of disorder that affect human beings
 c) Study the HIV disease development
 d) Developing genome project of other animals
117. In human beings 99.9% of genome sequence are same in all individuals only 0.1% of genome differ that **1**
 a) Make every individual similar in phenotypic appearance
 b) Make every individual genetically similar
 c) Make every individual unique in phenotypic appearance
 d) Make genetic variation for evolution
118. Gel electrophoresis is used for **1**
 a) Isolation of DNA molecule
 b) Separation of DNA fragments according to their size
 c) Cutting of DNA into fragments
 d) Construction of recombinant DNA by joining with cloning vectors
119. Repetitive DNA sequence shows high degree of polymorphism that forms the basis of **1**
 a) Cell division
 b) Cell differentiation
 c) DNA fingerprinting
 d) Genetic disease assessment
120. The technique of DNA fingerprinting was initially developed by **1**
 a) S. Mond
 b) Alec Jefferys
 c) Robert Sanford
 d) D.Pollard


Solution
Class 12 - Physics
MCQ Examination July (2019-20)
Section A

1. (a)

$$2 \frac{k\lambda}{R}$$

Explanation:

Consider a uniformly charged thin rod bent into a semicircle of radius R.



Charge per unit length: $\lambda = \frac{Q}{\pi R}$

Charge on slice: $dq = \lambda R d\theta$ (taken positive) Electric field generated by slice:

$$dE = \frac{k|dq|}{R^2} = \frac{k|\lambda|d\theta}{R} \text{ directed radially (inward for } \lambda > 0 \text{)}$$

Components of dE, $dE_x = dE \cos \theta$,

$$dE_y = -dE \sin \theta$$

Electric field from all slices added up: $E_x = \frac{k\lambda}{R} \int_0^\pi \cos \theta \, d\theta = \frac{k\lambda}{R} [\sin \pi - \sin 0] =$

0

$$E_y = -\frac{k\lambda}{R} \int_0^\pi \sin \theta \, d\theta = \frac{k\lambda}{R} [\cos \pi - \cos 0] = -\frac{2k\lambda}{R}$$

2. (a)

$$\frac{\sigma^2 R^2}{\epsilon_0}$$

Explanation:

Outward electric field at the surface of shell is $E = \frac{\sigma}{2\epsilon_0}$ If Q is the charge on the shell and A is the area,

than the outward pressure is $P = \frac{QE}{A} = \sigma E = \frac{\sigma^2}{2\epsilon_0}$

Force = PX effective area of hemispherical shell = $\frac{\sigma^2}{2\epsilon_0} \times \pi R^2$

So $F \propto \frac{\sigma^2}{\epsilon_0} R^2$

3. (b)

The magnitude of the force between the charges at C and B is $\frac{q^2}{54\pi\epsilon_0 R^2}$

Explanation:

The electric field due to charges at A and B are equal and opposite, So at O the electric field is due to C only, which has a magnitude

$$E = \frac{2q}{12\pi_0 R^2} = \frac{q}{6\pi_0 R^2}$$

The potential energy of the system is not zero. Potential at O is zero and Force between B and C

$$F = \frac{\frac{q}{3} \frac{2q}{3}}{4\pi_0 (2R \sin 60^\circ)^2} = \frac{q^2}{54\pi_0 R^2}$$

4. (b)

$$q = \frac{Q}{2}$$

Explanation:

Let, q and (Q-q) and 'r' be the separation between the charges.

The force of repulsion between them is,

$$F = \frac{K(Q-q)q}{r^2} = \frac{k}{r^2} (Qq - q^2)$$

Differentiation F w.r.t. q and setting it to zero will give us the extremum force.

$$\frac{dF}{dq} = \frac{k}{r^2} \frac{d}{dq} (Qq - q^2) = 0$$

$$\Rightarrow \frac{k}{r^2} (Q - 2q) = 0$$

$$\Rightarrow Q - 2q = 0$$

$$\Rightarrow q = \frac{Q}{2}$$

For this value of q, the force is extremum (minimum or maximum). The force will be maximum if the second differentiation of F is less than zero.

$$\frac{d^2 F}{dq^2} = \frac{-2k}{r^2} < 0$$

Thus, the force of repulsion is maximum when $q = \frac{Q}{2}$

5. (c)

$$1.45 \times 10^{-3} \text{C}, 1.6 \times 10^8 \text{Nm}^2/\text{C}$$

Explanation:

$$\text{a. } r = \frac{d}{2} = \frac{2.4}{2} = 1.2 \text{m}$$

$$\sigma = 80 \times 10^{-6} \text{C/m}^2$$

$$\sigma = \frac{q}{4\pi r^2}$$

$$80 \times 10^{-6} = \frac{q}{4 \times 3.14 \times (1.2)^2}$$

$$q = 1.45 \times 10^{-3} C$$

$$b. \phi = \frac{q}{\epsilon_0} = \frac{1.45 \times 10^{-3}}{8.85 \times 10^{-12}} = 1.6 \times 10^8 Nm^2/C$$

6. (b)

Zero

Explanation:

The field of opposite charges cancels each other so net electric field at centre = 0

7. (c)

Zero

Explanation:

On all the dipoles net charge = 0, hence net charge enclosed within the surface = 0. so the total electric flux coming out of the surface $\phi = \frac{q_{net}}{\epsilon_0} = 0$

8. (d)

Electrons flow from the conductor to the earth

Explanation:

After earthing a positively charged conductor electrons flow from earth to conductor and if a negatively charged conductor is earthed then electrons flows from conductor to earth.



9. (d)

The angular momentum of the charge $-q$ is constant

Explanation:

Since the charge $-q$ is moving in elliptical orbit so to make its motion stable the total angular momentum of the charge is constant since it experience a centripetal force from the charge $+Q$ so it follow the motion as the motion of earth around sun.

10. (b)

None of these

Explanation:

Electric flux for any closed surface is defined as $\oint \vec{E} \cdot \vec{ds}$.

The flux through ABCD can be calculated, by first taking a small elemental surface and then writing the $\vec{E} \cdot \vec{ds}$ for this element, keep in mind that electric field at the location of this element is the resultant of both the charges. It is quite obvious the flux through ABCD comes out to be non-zero because at every point of the surface, the angle between E and ds is less than 90° giving a positive non-zero value for the entire surface.

The dimension of flux should be that of $\frac{q}{\epsilon_0}$, where all given options have dimensional formula for $\frac{q}{\epsilon_0 l}$.

11. (a)

$30\text{Nm}^2/\text{C}$, $15\text{Nm}^2/\text{C}$

Explanation:

1. Electric field intensity, $= 3 \times 10^3 \hat{i} \text{ N/C}$

Magnitude of electric field intensity, $= 3 \times 10^3 \text{ N/C}$

Side of the square, $s = 10 \text{ cm} = 0.1 \text{ m}$

Area of the square, $A = s^2 = 0.01 \text{ m}^2$

The plane of the square is parallel to the y-z plane.

Hence, angle between the unit vector normal to the plane and electric field,

$\theta = 0^\circ$ Flux (Φ) through the plane is given by the relation, $\Phi =$

$$\vec{E} \cdot \vec{A} = EA \cos\theta = 3 \times 10^3 \times 0.01 \times \cos 0^\circ = 30 \text{ N m}^2/\text{C}$$

2. Electric field intensity, $= 3 \times 10^3 \hat{i} \text{ N/C}$

Magnitude of electric field intensity, $= 3 \times 10^3 \text{ N/C}$

Side of the square, $s = 10 \text{ cm} = 0.1 \text{ m}$

Area of the square, $A = s^2 = 0.01 \text{ m}^2$.

Angle between the unit vector normal to the plane and electric field, $\theta = 60^\circ$

Flux (Φ) through the plane is given by the relation, $\Phi = \vec{E} \cdot \vec{A} = EA \cos\theta =$

$$3 \times 10^3 \times 0.01 \times \cos 60^\circ = 15 \text{ N m}^2/\text{C}$$

12. (d)
 $0.07 \mu\text{C}$, No

Explanation:

a. Net outward flux through the surface of the box, $\phi = 8.0 \times 10^3 \text{ N m}^2/\text{C}$

For a body containing net charge q ,

flux is given by the relation, $\epsilon_0 = \text{Permittivity of free space} = 8.854 \times 10^{-12} \text{ N}^{-1} \text{C}^2 \text{ m}^{-2}$

We have

$\phi = \frac{q}{\epsilon_0}$ so $q = \epsilon_0 \phi = 8.854 \times 10^{-12} \times 8.0 \times 10^3 = 7.08 \times 10^{-8} = 0.07 \mu\text{C}$ Therefore, the net charge inside the box is $0.07 \mu\text{C}$.

b. No Net flux piercing out through a body depends on the net charge contained in the body. If net flux is zero, then it can be inferred that net charge inside the body is zero. The body may have equal amount of positive and negative charges.


13. (b)
 decrease

Explanation:

Since the spheres are conducting, the surface charge distribution on each sphere will be altered because of the repulsion from the charges on the other sphere. In particular, the charges on each sphere will be pushed away by the charges on the other sphere. This will cause the charges on opposite spheres to be further away from each other, and the force of repulsion to be less than in the case of a uniform surface charge distribution.

14. (c)
 $\frac{\lambda}{2\pi\epsilon_0 a}$

Explanation:



I have used the symbol R for radius in the diagram.

Let λ be the linear charge density .then a small charge element $dq = \lambda a d\phi$ and

electric field due to this element at centre of arc $dE = \frac{dq}{4\pi\epsilon_0 a^2}$

For every dq there exist a dq' such that y component of dE cancels out thus

$$E_x = \int_{-\pi/2}^{\pi/2} dE \cos \phi. \text{ Substitute for } dE \text{ and } dq$$

$$E_x = \int_{-\pi/2}^{\pi/2} \frac{\lambda a \cos \phi d\phi}{4\pi \epsilon_0 a^2} \text{ on solving integral.}$$

$$E_x = \frac{\lambda}{2\pi\epsilon_0 a}$$

15. (d)


2C

Explanation:

The capacitance C of a parallel plate capacitor is given by $C = \frac{\epsilon_0 A}{d}$

A metal plate of thickness $d/2$ when introduced between the plates reduces the distance between the plates to $\frac{d}{2}$. The effective capacitance becomes

$$C_m = \frac{\epsilon_0 A}{\frac{d}{2}} = \frac{2\epsilon_0 A}{d} = 2C$$



Another explanation: The system can be considered to be three capacitors C_1 , C_2 , and C_3 connected in series.

$$C_1 = \frac{\epsilon_0 A}{x}; C_2 = \frac{\epsilon_0 K A}{\frac{d}{2}}; C_3 = \frac{\epsilon_0 A}{\frac{d}{2} - x}$$

K of a metal is infinity. $C_2 = \infty$. The equivalent capacitance

$$\frac{1}{C_m} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} = \frac{x}{\epsilon_0 A} + \frac{1}{\infty} + \frac{\frac{d}{2} - x}{\epsilon_0 A}$$

$$= \frac{1}{\epsilon_0 A} \left[x + \frac{d}{2} - x \right] = \frac{\frac{d}{2}}{\epsilon_0 A}$$

$$C_m = \frac{2\epsilon_0 A}{d} = 2C$$

16. (b)

32.0

Explanation:

Each capacitor of capacitance $8\mu F$ can withstand a maximum potential of 250 V.

When equal capacitors are connected in series, the potential difference across them is equal.

If there are m capacitors in series such that the potential across each is 250 V, then, $\frac{1000}{m} = 250; m = 4$.

The equivalent capacitance of 4 capacitors connected in series is

$$C_S = \frac{C}{m} = \frac{8}{4} = 2\mu F .$$

To achieve a capacitance of 16, n such rows of capacitors need to be connected in parallel.

$$C_{eq} = nC_S = 16\mu F; n = \frac{16}{C_S} = \frac{16}{2} = 8 .$$

To make a condenser of $16\mu F$, 8 rows of capacitors with each row containing 4 capacitors are to be connected.

The total number of capacitors = $n \times m = 4 \times 8 = 32$.

17. (a)

$$\frac{1}{4}C(V_1 - V_2)^2$$

Explanation:

The initial energy of the two capacitors $U_i = \frac{1}{2}CV_1^2 + \frac{1}{2}CV_2^2$.

The charges on the capacitors are $Q_1 = CV_1; Q_2 = CV_2$

When they are joined, they attain a common potential V . $V = \frac{\text{total charge}}{\text{total capacitance}}$

$$= \frac{Q_1 + Q_2}{C + C} = \frac{CV_1 + CV_2}{2C} = \frac{V_1 + V_2}{2} .$$

Final energy $U_f = \frac{1}{2}CV^2 + \frac{1}{2}CV^2 = CV^2$ Loss of energy

$$U_i - U_f = \frac{1}{2}C(V_1^2 + V_2^2) - CV^2$$

$$= \frac{1}{2}C(V_1^2 + V_2^2) - C\left(\frac{V_1 + V_2}{2}\right)^2$$

$$= \frac{1}{4}C(V_1 - V_2)^2$$

18. (b)

$$0.17 \times 10^{-2} \text{ J}$$

Explanation:

Explanation here: (A) Initial energy

$$U_1 = \frac{1}{2}C_1V_1^2 + \frac{1}{2}C_2V_2^2$$

$$= \frac{1}{2}(2 \times 10^{-6})(100)^2 + \frac{1}{2}(4 \times 10^{-6})(50)^2$$

$$= 1.5 \times 10^{-2} J$$

The common potential after they are connected in parallel

$$V = \frac{C_1 V_1 + C_2 V_2}{C_1 + C_2}$$

$$= \frac{(2 \times 10^{-6})(100) + (4 \times 10^{-6})(50)}{(2 \times 10^{-6}) + (4 \times 10^{-6})}$$

$$= \frac{2}{3} \times 10^2 V$$

The final energy

$$U_2 = \frac{1}{2} (C_1 + C_2) V^2$$

$$= \frac{1}{2} [(2 \times 10^{-6}) + (4 \times 10^{-6})] \left(\frac{2}{3} \times 10^2 \right)^2$$


$$= 1.33 \times 10^{-2} J .$$

So change in energy is $\Delta U = U_1 - U_2 = 0.17 \times 10^{-2}$

19. (b)

$9 \mu F$

Explanation:



If P is at positive potential, then Q is at negative potential and R is at positive potential. The system therefore reduces to 3 capacitors in parallel. $C = 9 \mu F$

20. (a) Decreasing the battery potential

Explanation:

An electroscopes is a device which measures the potential difference. If it is connected in parallel to the capacitor, the potential across it will be equal to the potential across the capacitor, which is equal to the potential across the battery. On decreasing the battery potential, the potential difference across the electroscopes reduces and hence the reading reduces. While the capacitor is connected to the battery, Placing a dielectric between the plates, or decreasing the distance between the plates or increasing the area of the plates will not change the potential difference across it; since it will always remain equal to the potential difference maintained by the battery. In the cases B, C and D, The capacitance of the capacitor, however increases; but this increase happens due to increase in the charge stored in the capacitor while the potential remains constant.

21. (d)
 $3 \mu\text{F}$

Explanation:

The capacitance of the first capacitor $C = \frac{\epsilon_0 A}{d} = 2 \mu\text{F}$ The second capacitor is considered to be made of two capacitors C_1 (air filled) and C_2 (dielectric) connected in series.

$$C_1 = \frac{\epsilon_0 A}{\frac{d}{2}} = 2C = 4 \mu\text{F};$$

$$C_2 = \frac{K \epsilon_0 A}{\frac{d}{2}} = 2KC = 12 \mu\text{F}$$

The equivalent capacitance

$$C_1 = \frac{\epsilon_0 A}{\frac{d}{2}} = 2C = 4 \mu\text{F};$$

$$C_2 = \frac{K \epsilon_0 A}{\frac{d}{2}} = 2KC = 12 \mu\text{F}$$

22. (d)

the potential difference across the plates of A is 6 V and across the plates of B is 4 V

Explanation:

the potential difference across the plates of A is 6 V and across the plates of B is 4 V

23. (a)

$$d = 10^{-5} \text{m}, A = 10^{-2} \text{m}^2$$

Explanation:

The capacitance of a parallel plate capacitor of area A, plate separation d with a dielectric of dielectric constant K is $C = \frac{\epsilon_0 K A}{d}$.

$$\text{The ratio } \frac{A}{d} = \frac{C}{\epsilon_0 K} = \frac{1.77 \times 10^{-6}}{8.85 \times 10^{-12} \times 200} = 10^3 .$$

The minimum plate separation d' for which the capacitor will not breakdown is found using $E = \frac{V}{d'}$

where E is the breakdown strength and V is the maximum potential the capacitor can withstand.

$$d' = \frac{V}{E} = \frac{20}{3 \times 10^6} = 6.67 \times 10^{-6} \text{m} .$$

The plate separation has to be greater than $6.67 \times 10^{-6} m$
 therefore only option A satisfies the condition $\frac{A}{d} = \frac{10^{-2}}{10^{-5}} = 10^3$

24. (c)
 $4 \mu F$

Explanation:

The capacitance of the first capacitor $C = \frac{\epsilon_0 A}{d} = 2 \mu F$

The second capacitor is considered to be made of two capacitors C_1 (air filled) and C_2 (dielectric) connected in parallel.

$$C_1 = \frac{\epsilon_0 \frac{A}{2}}{d} = \frac{C}{2} = 1 \mu F; C_2 = \frac{K \epsilon_0 \frac{A}{2}}{d} = \frac{KC}{2} = 3 \mu F$$

$$C_{eq} = C_1 + C_2 = 1 \mu F + 3 \mu F = 4 \mu F$$

25. (d)
 $\frac{q^2}{2\epsilon_0 A}$

Explanation:

Force between two plates of the capacitor

$F = uA$ where u is

The energy density $u = \frac{1}{2} \epsilon_0 E^2$

The electric field $E = \frac{\sigma}{\epsilon_0}$

and the charge density $\sigma = \frac{q}{A}$

$$F = \frac{1}{2} \epsilon_0 E^2 A = \frac{1}{2} \epsilon_0 \left(\frac{\sigma}{\epsilon_0} \right)^2 A = \frac{1}{2} \frac{\sigma^2 A}{\epsilon_0} = \frac{1}{2} \left(\frac{q}{A} \right)^2 \frac{A}{\epsilon_0} = \frac{q^2}{2A\epsilon_0}$$

26. (c)
 10%

Explanation:

Let original Current In lamp = I

Resistance of Lamp = R

Then power $P = I^2 R$

According to question,

$$\text{New Current } I_n = I - I \times \frac{5}{100} = \frac{19}{20} I$$

Resistance = R

$$\text{New power } P_n = I_n^2 R = \left(\frac{19}{20} I \right)^2 R = \frac{361}{400} I^2 R$$

$$\text{Power decrease} = I^2 R - \frac{361}{400} I^2 R = \frac{39}{400} I^2 R$$

$$\% \text{ Decrease} = \frac{\text{change in power}}{\text{original power}} \times 100$$

$$= \frac{\frac{39}{400} I^2 R}{I^2 R} \times 100 = \frac{39 I^2 R}{400 I^2 R} \times 100$$

$$= \frac{39}{4} = 9.75\% \approx 10\%$$

27. (d)

5 mV/m

Explanation:

The total resistance is the sum of the resistance of the potentiometer and the external resistance.

$$R = R_{\text{pot}} + R_{\text{ext}} = 5 + 995 = 1000 \text{ ohms .}$$

The current through the potentiometer wire $I = \frac{E}{R} = \frac{10}{1000} = 0.01 \text{ A}$ $I = E/R = 10/1000 = 0.01 \text{ A}$.

The potential drop across the potentiometer wire is

$$V = I \times R_{\text{pot}}$$

$$\Rightarrow V = 0.01 \times 5$$

$$V = 0.05 \text{ V}$$

The potential gradient = (potential drop across the potentiometer wire)/ length of the potentiometer wire)

$$= \frac{0.05}{10}$$

$$= 5 \times 10^{-3} \text{ V/m}$$

$$= 5 \text{ mV/m}$$

28. (a)

$$I^2 R$$

Explanation:

The power dissipated

$$P = V \times I$$

$$\text{Since } V = IR$$

$$P = I^2 R$$

29. (a)

The electric current I flowing through a substance is proportional to the voltage V across its ends

Explanation:

Ohm's law states I is proportional to V. This holds good at steady temperatures and for the flow of constant current.

30. (a)
93.0%

Explanation:

$V = 200 \text{ Volt}$ $I = 3 \text{ A}$ $\text{time} = 10 \text{ minute} = 600 \text{ sec}$

The electric energy (input energy) = $VIt = 200 \times 3 \times 600 = 360000 \text{ joule}$

$m = 1 \text{ l} = 1000 \text{ g} = 1 \text{ kg}$ sp. heat of water = $4186 \text{ j/kg}^{\circ}\text{C}$ $\Delta T = 100 - 20 = 80^{\circ}\text{C}$

Heat energy (output energy) = $mc\Delta T = 1 \times 4186 \times 80 = 334880 \text{ joule}$

efficiency = $= \frac{\text{output energy}}{\text{input energy}} \times 100 = \frac{334880}{360000} \times 100 = 93.0\%$

31. (d)

Increase in the rate of collisions between the carriers and vibrating atoms of the conductor

Explanation:

When temperature increases, the thermal speed of the electrons increases as well as, the amplitude of vibration of the positive ions inside the metal conductor also increase, about their mean positions. Thus, the collisions between the electrons and the positive metal ions become more frequent and this decreases the relaxation time, t , leading to an increase in the resistivity of the conductor.

32. (c)
 $\frac{\text{Volt meter}}{\text{Ampere}}$

Explanation:

\therefore Resistance $R = \rho \frac{L}{A}$

Where ρ is resistivity, L is length and A is area.

$\Rightarrow \rho = R \frac{A}{L}$

also $R = \frac{V}{I}$

$\therefore \rho = \frac{V \times A}{I \times L}$

and in units,

$$\rho = \frac{(\text{Volts}) \times (\text{meter})^2}{(\text{Ampere}) \times (\text{meter})}$$

$$\Rightarrow \rho = \frac{\text{Volt meter}}{\text{Ampere}}$$

33. (a)

0.4 A

Explanation:

If the battery has an e.m.f E , resistance of the potentiometer is R and the internal resistance of the battery is r , then the current I flowing in the potentiometer wire is given as

$$I = \frac{E}{(R+r)}$$

$$I = \frac{2}{(4+1)}$$

$$I = 0.4 \text{ A}$$

34. (c)

the net current flowing through the area normally per unit time

Explanation:

Current density $J = I/A$

In electromagnetism, current density is the electric current per unit area of cross section. It is a vector and has a direction along the area vector.

35. (b)

is the direction in which positive charges move

Explanation:

Current flows in a conductor due to the flow of negatively charged electrons. However, the direction of conventional current is taken to be opposite to the direction of flow of electrons. It can therefore be considered as the direction in which positive charges move.

36. (d)

$$10^6 \text{ m/s}, 10^{-4} \text{ m/s}$$

Explanation:

The random velocities of electrons is of the order 10^5 to 10^6 m/s, while the drift velocities are of the order 0.1mm/s (10^{-4} m/s)

37. (b)

$$\frac{1}{3}\Omega$$

Explanation:

Given,

$$E_1 = E_2 = 1.5$$

Let internal resistance of battery be r . If batteries are connected in series then,

$$E = E_1 + E_2 = 3 V$$

$$T_{\text{total}} = 2r$$

Now,

$$I = \frac{E}{(R+2r)}$$

$$\Rightarrow 1 = \frac{3}{(R+2r)}$$

$$R + 2r = 3 \dots (i)$$

If batteries are connected in parallel

$$E = 1.5V$$

$$\frac{1}{T_{\text{total}}} = \frac{1}{r} + \frac{1}{r}$$

$$T_{\text{total}} = \frac{r}{2}$$

and,

$$0.6 = \frac{1.5}{\left(R + \frac{r}{2}\right)}$$

$$\Rightarrow 0.6R + 0.3r = 1.5 \dots (ii)$$

on solving equation (i) and (ii)

$$r = \frac{1}{3}\Omega$$

38. (b)

The algebraic sum of changes in potential around any closed loop must be zero.

Explanation:


Kirchhoff's loop rule is based on the principle of conservation of energy. Since work done in transporting a charge in a closed loop is zero. The algebraic sum (since potential differences can be both positive and negative) of potential differences around any closed loop is always zero.

39. (a)

0.2 A current flows in G.

Explanation:

In steady state, no current flows through the capacitors.



(B) The current flows along ABGDCA. The resistances 4Ω , 1Ω and 5Ω are in series. Total resistance of the circuit = $R = 4 + 1 + 5 = 10\Omega$. Current $I = V/R = 2/10 = 0.2\text{ A}$. The current through the galvanometer is 0.2 A

40. (b)

nIA

Explanation:

Magnetic moment is defined as the product of total current and area of loop

$$M = n \times I \times A$$

Solution
Class 12 - Chemistry
MCQ July
Section A

41. (a)
2

Explanation:

As initial concentration is increased half life is decreasing so order of reaction is 2.

for second order reaction, $rate \propto \frac{1}{[R]}$

42. (c)
More than ΔH

Explanation:

$\Delta H = +ve$ for endothermic reaction
, therefore, $E_a > \Delta H$

43. (d)
Zero order

Explanation:

Uniform Rate of reaction is independent of concentration of reactants.

44. (b)
Concentration of reactants keep on changing

Explanation:

Rate of reaction is dependent on concentration of reactants. if concentration of reactants change then rate of reaction become non-uniform.

45. (d)
is halved by half by reducing the concentration of RCl

Explanation:

since rate of reaction = $k[\text{RCl}]^1$

so if conc. of RCl is halved the rate of reaction will also become half.

46. (c)

1200 min

Explanation:

$$t_{99.9} = 10 \times t_{1/2}$$

detail:

$$\text{here, } k = \frac{0.693}{120}$$

$$\text{also, } t = \frac{2.303 \times 120}{0.693} \log 10^3 = \frac{2.303 \times 120 \times 3}{0.693} \log 10$$

$$\Rightarrow t = \frac{2.303 \times 120 \times 3 \times 1}{0.693} = 1196.36 \simeq 1200$$

47. (c)

104 kJ/mol

Explanation:

$$\ln K = \ln A - \frac{E_a}{RT}$$

on comparing with $y = mx + c$

$$\text{slope} = \frac{E_a}{2.303R}$$

$$E_a = 5.42 \times 10^3 \times 2.303 \times 8.314$$

$$E_a = 103.7 \text{ KJ/mol}$$

48. (b)

x + y

Explanation:

Order of reaction with respect to A is x and w.r.t to B is y so total order of reaction is x+y.

49. (b)

Rate constant depend upon the concentration of the reactants


Explanation:

Rate constant is independent of concentration of reactant for a particular reaction.

50. (b)

8 times

Explanation:



$$\Rightarrow Rate = Rate_2 = k_2[NO][NOBr_2]$$

$$\rightarrow Rate_1 = Rate_{-1} \rightarrow k_1[NO][Br_2] = k_{-1}[NOBr_2]$$

$$\rightarrow [NOBr_2] = (k_1/k_{-1})[NO][Br_2]$$

$$\Rightarrow Rate = k_2[NO][NOBr_2] = k_2[NO](k_1/k_{-1})[NO][Br_2]$$

$$\Rightarrow Rate = (k_2k_1/k_{-1})[NO]^2[Br_2] = k[NO]^2[Br_2]$$

Rate = $k[NO]^2[Br_2]$, since rate of reaction w.r.t [NO] is second order and w.r.t [Br] is first order, then rate of reaction become 8 times when conc. of [NO] and [Br] is doubled.

$$rate' = k[2NO]^2 [2Br_2]$$

$$rate' = 8 \times Rate$$

51. (d)

$$mol^{-1} litre s^{-1}$$

Explanation:

unit of rate constant for nth order of reaction are:

$$\text{unit of } k \text{ for nth order} = (molL^{-1})^{1-n} s^{-1}$$

put n=2 for second order reaction.

52. (a)

Elementary reaction


Explanation:

An elementary reaction is a chemical reaction in which one or more chemical species react directly to form products in a single reaction step and with a single transition state

53. (b)

$$0.1 molL^{-1}min^{-1}$$

Explanation:



$$\text{rate} = -\frac{1}{2} \frac{d[X]}{dt} = -\frac{d[Y]}{dt} = \frac{d[Z]}{dt}$$

$$\frac{d[X]}{dt} = 2 \frac{d[Z]}{dt} = 2 \times 0.05 = 0.1 \text{ mol L}^{-1} \text{ min}^{-1}$$

54. (b)

$$\text{rate} = K [A] [B]^2$$

Explanation:

$$\text{rate} = K [A] [B]^2$$

since rate of given reaction is first order wrt A reactant and second order wrt B reactant.

order of reaction is sum of powers of each reactant in rate law expression.


so, order of reaction = 1 + 2 = 3

55. (b)

Iron

Explanation:


Finely divided Iron (Fe) is used as a catalyst as the surface area of small particles is much larger than normal crystal. Along with Fe a promoter (substance that activates a catalyst) Molybdenum (Mo) is used.



According to Le Chatelier's principle, high pressure and temperature promote this reaction in forward direction.

Also, Iron oxide (Fe₂O₃) along with potassium oxide and alumina is used for Haber's process.

56. (b)



Explanation:

It is a redox reaction where sulphur is oxidized and nitrogen is reduced and result will be a colloidal solution.

57. (d)

Oxidation of oxalic acid by acidified KMnO₄

Explanation:

Autocatalysis occurs when the product of a reaction serves as a catalyst for the reaction.

58. (c)

Physical

Explanation:

Physical adsorption is favoured at low temperature because it involves only vanderwall interactions between adsorbate and adsorbent.

59. (d)

Due to Tyndall effect

Explanation:

This is because of tyndall effect caused by the scattering of light by colloidal particles of As_2S_3 .

60. (b)

Low temperature, high pressure

Explanation:

Physisorption is favoured only at low temperature and high pressure.

61. (c)

Alum

Explanation:

Alum is used in dying industries.

62. (c)

Mechanical

Explanation:


Movement is always a mechanical property.

63. (b)

Associated colloids

Explanation:

Micelles are chemical structures formed with both hydrophilic (they'll mix into water) and hydrophobic (they don't mix into water). Also called as Associated colloids. In the general case, micelles are formed when there is an ideal temperature in the medium (called the Kraft temperature) and a certain concentration of electrolytes (called the CMC: Critical Micelle Concentration) in the medium.



- i. Grease or oil on surface of cloth.
- ii. Stearate ions arranged around the grease or oil droplet.
- iii. Grease or oil droplet surrounded by stearate ions (ionic micelle formed).

64. (a)

Liquid in gas

Explanation:

Dispersed phase is liquid, dispersion medium is gas.

65. (b)

Flocculation

Explanation:

Due to very less size of colloidal solutions, they do not exhibit flocculation. When a sol is colloiddally unstable then the formation of aggregates is called flocculation.

66. (b)

Vanadium pentoxide

Explanation:

V_2O_5 is used as catalyst in contact process .

67. (a)

Thermite process

Explanation:

Thermite process doesn't require a catalyst. It can easily proceed without the help of catalyst.

68. (d)
Combustion zone

Explanation:

Combustion zone maintains the highest temperature around 1775 K.
e.g. extraction of Fe

69. (d)
Leaching

Explanation:

Leaching is a process in which ore is digested with a solvent to form a soluble complex.

Example: Leaching of alumina from bauxite.

70. (a)
Ag

Explanation:

Ag is obtained by Leaching process by using dil. NaCN/KCN followed by replacement to give the pure metal.

71. (c)
 Al_2O_3 and Na_3AlF_6

Explanation:

Al_2O_3 and Na_3AlF_6 (molten solution). Aluminium oxide has a very high melting point (over 2,000°C), so it would be expensive to melt it. Instead, it is dissolved in molten cryolite, an aluminium compound with a lower melting point than aluminium oxide. The use of cryolite reduces some of the energy costs involved in extracting aluminium.

72. (d)
3%

Explanation:

Cast iron is made by melting pig iron with scrap iron and coke using hot air blast. It has 3% of carbon content and is extremely hard and brittle.

73. (a)

Sn

Explanation:


Cassiterite is a tin dioxide mineral. It is generally opaque, but it is translucent in thin crystals. Its luster and multiple crystal faces produce a desirable gem. Cassiterite has been the chief tin ore throughout ancient history and remains the most important source of tin today.

74. (c)

$\text{Cu} + \text{SO}_2$

Explanation:

This auto reduction reaction gives metallic copper and sulphur dioxide.



75. (b)

Electrolytic refining

Explanation:

In this method, the impure metal acts as anode. A strip of same pure metal is used as cathode. A salt of metal is made an electrolyte. On passing electricity through the solution, the pure metal moves towards the cathode, and impurities present in the anode settle down at the bottom as anode mud.

76. (c)

Mercury

Explanation:

HgS is brick red form of sulphide ore of Hg from which it can be profitably extracted. It resembles quartz in symmetry.

77. (b)

Hydraulic washing

Explanation:

This is hydraulic washing or gravity separation. Here when stream of water is passed it takes away all the lighter impurities with it and the heavier ore particles are left behind.

78. (d)

Aniline

Explanation:



During froth flotation, substances are used to stabilize the froth so that it can be easily skimmed off and purified. Aniline and cresols are froth stabilizers.

79. (d)

Carbon monoxide

Explanation:

CO is used as reducing agent in blast furnace to get iron at such a high temperature.



80. (a)

Iron

Explanation:

Iron is 2nd most abundant metal in earth's crust around 5 %.

Solution
Class 12 - Biology
MCQ July 2019
Section A

81. (d)

Haemophilia and red green colour blindness

Explanation:

Haemophilia and red-green colour blindness both are a sex-linked recessive gene on X chromosome. Body height is an example of polygenic inheritance. Rhesus blood group is based on the presence or absence of Rh-protein on the surface of RBC, phenylketonuria (PKU) is a recessive autosomal variation.

82. (b)

Down's syndrome

Explanation:

Mendelian disorders are mainly determined by alteration or mutation in single gene. These disorders are transmitted to the offspring on the same line as principles of inheritance. Down's syndrome is due to presence of additional copy of chromosome number 21.

83. (b)

Haemophilia

Explanation:

The inheritance pattern of a certain type of trait in humans shown above is haemophilia.

A son cannot inherit the defective gene from his father. This is a recessive trait and can be passed on if cases are more severe with the carrier. Genetic testing and genetic counselling are recommended for families with haemophilia. The disease is X-linked and the father cannot pass haemophilia through the Y-chromosome.

84. (d)

Homologous pairs

Explanation:

Somatic cells of human beings contain 23 pairs of chromosomes, 23 each from male and female gametes. These chromosomes always occur as homologous chromosomes as they contain the same kinds of traits.

85. (a)

Hardly any cross-overs are produced

Explanation:

When two genes are situated very close to one another on a chromosome, hardly any cross-overs are produced. Such genes are called linkage genes and do not separate from each other during gamete formation.

86. (b)

All statements are correct

Explanation:

Physical association between two genes located on the same chromosome is called linkage. The generation of non-parental gene combinations is called recombination. Recombination is essential for the generation of new traits in a species.

87. (c)

Acrocentric

Explanation:

A chromosome with a sub-terminal centromere near the middle is called an acrocentric chromosome. In an acrocentric chromosome, one arm is larger and the other is shorter.

88. (c)

Complementary genes

Explanation:

Complementary genes are one of two or more genes that when present together produce effects qualitatively distinct from the separate effect of any one of them.

89. (a)

An increase in whole set of chromosome in organism

Explanation:

Cytokinesis is the division of chromosome. Failure of cytokinesis after telophase stage of cell division results in the an increase in whole set of chromosome in organism that leads to polyploidy.

90. (b)

Euploidy

Explanation:

Most of the organisms are diploid, contain two set of chromosome ($2n$) but a number of organisms contains multiple of chromosome set like $3n, 4n, 6n$ etc. This condition of having multiple sets chromosome is called Euploidy.

91. (c)

A several generation

Explanation:

Pedigree analysis is the study of particular traits in several generation of a family. In this analysis, inheritance of particular trait is represented in family tree over the generation.

92. (b)

Male Drosophila

Explanation:

Karyotype is the representation of a chromosome in order of shape and size. Male Drosophila fly have 3 pairs of autosomes and 1 pair (XY) sex chromosome.

93. (d)

X-chromosome

Explanation:

Henking trace a specific structure in 50% of sperms. He gave the name x-body to this structure but he was not able to explain the significance of this

structure. Further, investigation revealed that x-body was a chromosome, called X-chromosome.

94. (c)

Cross over percentage

Explanation:

Map distance is tools to find the distance between two genes on a chromosome. The genes closer to each other shows less number of crossing over percentage.

95. (d)

Klinefelter's syndrome - 44 autosomes +XXY

Explanation:

Klinefelter's syndrome is due to presence of 44 autosomes +XXY sex chromosomes. Color blindness is X-linked recessive trait, Erythroblastosis foetalis is due to O-negative blood group and down's syndrome is due to additional copy of chromosome number 21.

96. (a)

XX - XO type of sex determination

Explanation:

The chart given below represents XX-XO type of sex determination in which XX forms the female and XO develops as a male. It is common in birds.

97. (b)

Segregation

Explanation:

Linkage is the non-separation of two genes present on same chromosome. Segregation is the separation of two alleles from each other. So segregation is opposite to linkage.

98. (c)

This disease is due to a X-linked recessive mutation

Explanation:

Hemophilia is sex-linked recessive disease due to mutation in X-chromosome. In this disease blood clotting do not takes place in case of injury and minor cut leads to death of individual.

99. (c)

All colour blind

Explanation:

Colour blindness is sex-linked genetic disorder in which person is not able to detect the colour of the object. Red and green colour cannot be distinguished by them. It is carried by X-chromosome. Color blind woman transfer their X-chromosome to all offspring but daughter will be carrier and son will be all colour blind.

100. (c)

incorrect

Explanation:

Women are not responsible for sex of child because they produce only one kind of gamete containing X-sex chromosome, on the other hand male produce two kinds of sperms half containing X and half Y sex chromosome.

101. (b)

Statement I and II is correct

Explanation:

DNA finger printing is highly reliable method of identification of individual involved in crimes. DNA a fingerprinting is a sure method in solving paternity and maternity disputes. DNA fingerprinting cannot be used to cure HIV infection.

102. (c)

Monozygotic twins or identical twins

Explanation:

Autoradiogram of VNTR probe gives many band of different size. It differs from individual to individual except monozygotic twins or identical twins Because MZ twins share the same genetic makeup (DNA) because they are formed from a single zygote (fertilized egg).

103. (d)

Cistron

Explanation:

In biochemical genetics the term gene is being replaced by cistron. Cistron is a segment of DNA consisting of a stretch of deoxyribonucleotides which code for a biochemical controlling other cistron.

104. (d)

Polymerase chain reaction (PCR) by amplification process

Explanation:

Amplification is a mechanism leading to multiple copies of a chromosomal region within a chromosome arm.

The DNA amplification technique of the polymerase chain reaction (PCR) is a laboratory method for creating multiple copies of small segments of DNA.

105. (c)

3

Explanation:

In DNA molecules nitrogenous base of complementary strands binds with hydrogen bonds. In cytosine and guanine there are 3 hydrogen bonds and in adenine and thymine, the number of hydrogen bond is 2.

106. (a)

Cytoplasm

Explanation:

Transcription is the process of copying DNA code into mRNA. It occurs in the nucleus of the cell. The mRNA comes out of nucleus then its translation occurs into cytoplasm to form protein.

107. (b)

Released polypeptide chain

Explanation:

The figure shown above represents the translation process in which protein is produced. Ribosome provides the site for protein synthesis and t-RNA brings

the amino acids. The 'x' is the polypeptide chain produced.

108. (d)

Recombinant DNA techniques

Explanation:

Recombinant DNA technology, joining together of DNA molecules from two different species that are inserted into a host organism to produce new genetic combinations that are of value to science, medicine, agriculture, and industry. This technique made it possible to genetically engineer the genome of living organisms.

It involves a number of procedures like identification, separation, cloning and introducing into suitable vector.

109. (c)

Disease-associated sequences and tracing human history

Explanation:

Scientists have identified about 1.4 million locations where single base DNA differences (SNPs) occur in human. This information is helpful in finding chromosomal locations for disease-associated sequences and human history.

110. (a)

Splicing

Explanation:

The mRNA produced by transcription of DNA consists of exons and introns. The removal of introns and joining of exons to obtain mature mRNA is called splicing. It is followed by capping and tailing.

111. (d)

To identify the sequence of human DNA

Explanation:

The Human Genome Project was an international research effort to provide a complete and accurate sequence of the 3 billion DNA base pairs that make up the human genome and to find all of the estimated 20,000 to 25,000 human genes.

The Project was coordinated by the National Institutes of Health and the U.S. Department of Energy.

112. (b)

Regulator gene

Explanation:

Regulator gene is a gene that regulates the expression of one or more structural genes by controlling the production of a protein (such as a genetic repressor) which regulates their rate of transcription. Regulation of lac operon by repressor is called negative regulation.

113. (d)

Regulator gene

Explanation:

A regulator gene is a gene that codes for a repressor protein that inhibits the activity of an operator gene (a gene which binds repressor proteins thus inhibiting the translation of RNA to protein via RNA polymerase). In prokaryotes, regulator genes often code for repressor proteins.

114. (b)

Lactose

Explanation:

The lac operon (lactose operon) is an operon required for the transport and metabolism of lactose in *Escherichia coli* and many other enteric bacteria. Although glucose is the preferred carbon source for most bacteria, the lac operon allows for the effective digestion of lactose when glucose is not available.

115. (d)

DNA segments having radioactive isotopes

Explanation:

In finger printing the nylon membrane was incubated with radioactive probes. DNA Probes are small fragments of minisatellite DNA tagged with radioactive phosphorous. The probes only attach to the pieces of DNA that they are

complementary to – in this case they attach to the minisatellites in the genome. Radioactive isotopes can be easily identified using scanner.

116. (b)

Diagnose, treat and prevents the thousands of disorder that affect human beings

Explanation:

The Human Genome Project (HGP) is an international thirteen-year project that began on October 1990. It is important because it uses information from DNA to develop new ways to diagnose, treat, cure, or even prevent the thousands of diseases that afflict humankind.

117. (c)

Make every individual unique in phenotypic appearance

Explanation:

Genome variations are differences in the sequence of DNA from one person to the next

In human's 99.9% of the base sequences of DNA are same & are referred as **Bulk genomic DNA**.

The difference lies in remaining 0.1%. It is these differences which make every individual unique in their phenotypic appearance. This DNA has small stretches of **repetitive sequences**. They are referred as Repetitive DNA.

118. (b)

Separation of DNA fragments according to their size

Explanation:

Gel electrophoresis is used to separate macromolecules like DNA, RNA and proteins. DNA fragments are separated according to their size and proteins can be separated according to their size and their charge (different proteins have different charges) to study the genome of individual organism.

119. (c)

DNA fingerprinting

Explanation:

The term DNA fingerprinting - or genetic fingerprinting - is applied to the scientific process whereby samples of DNA are collected, collated and used to match other samples of DNA, which may have been found at the scene of a crime.

DNA fingerprinting works on the basis that each individual's DNA structure - or genetic make-up - is unique and therefore cannot be forged, faked or altered in any way.

One of the most common DNA fingerprinting procedures is RFLP: Restriction Fragment Length Polymorphism. Special enzymes are used to cut segments of a sample from which DNA is extracted. The procedure of RFLP focuses on repetitious sequences of DNA 'bases' which vary greatly from individual to individual.

120. (b)

Alec Jefferys

Explanation:

Sir Alec John Jeffreys, CH FRS (born 9 January 1950 in Oxford, Oxfordshire, England) is a British geneticist, who developed techniques for DNA fingerprinting and DNA profiling which are now used worldwide in forensic science to assist police detective work and to resolve paternity and immigration disputes.

He is a professor of genetics at the University of Leicester, and he became an honorary freeman of the City of Leicester on 26 November 1992.

In 1994, he was knighted for services to genetics